

Collective Flow vs. Hard Processes in Relativistic Heavy Ion Collisions

J. Burward-Hoy and B.V. Jacak

Department of Physics and Astronomy, SUNY Stony Brook, Stony Brook, NY 11794

In relativistic heavy ion collisions, the p_T distributions of hadrons reflect the transverse motion generated during the collision and therefore the collision dynamics. The moderate p_T values measured to date at $\sqrt{s} \approx 20$ GeV/nucleon contain a mix of hard and soft processes. The available data from the SPS on π , K, p, and Λ are used to constrain the magnitude of collective radial expansion. In $A + A$ collisions, the hydrodynamic distributions determined from the measured collective expansion are determined up to high p_T and are consistent with the π^0 data. To cleanly observe hard processes, measurements at $p_T \geq 4$ GeV/c will be required. Pions produced in $p + W$ collisions clearly deviate from hydrodynamic distributions at $p_T > 2$ GeV/c, underscoring the differences in the physics between $p + A$ and $A + A$ collisions.

PACS numbers: PACS numbers: 25.75.+r

The possibility of creating a new form of matter, a quark-gluon plasma, in ultra-relativistic heavy-ion collisions has generated intense experimental and theoretical effort. Transverse momentum distributions of hadrons reflect the transverse motion generated during the collision and therefore the collision dynamics. If there are sufficient secondary scatterings, local thermal equilibrium and collective hydrodynamical motion may occur. Measured hadronic transverse momentum (p_T) distributions have been used to search for such collective behavior and extract the hydrodynamic parameters [1, 2].

It is, however, the early stage of the collision, when the deconfined phase should exist, that is of paramount interest. Hard processes create probes for this phase, during the initial collisions between the participating nucleons. For example, particle jets arise from the fragmentation of high p_T quarks and gluons produced in initial hard scatterings. Study of high p_T hadrons can therefore yield information about energy loss of these quarks and gluons as they traverse the hot, dense nuclear medium. Theoretically, hard processes may be investigated using perturbative QCD-inspired parton models, tuned to reproduce experimental data on nucleon-nucleon (p+p) and nucleon-nucleus (p+A) collision systems [3]. Experimentally, the challenge is to identify a p_T range accessible with good statistical precision, but dominated by jets.

Transverse collective expansion of the colliding system has been observed in high energy nucleus-nucleus (A+A) collisions [1, 2]. This flow boosts the p_T of the hadrons, flattening the spectrum and effectively increasing the p_T range where soft processes are important. The π^0 production in Pb + Pb collisions at 158 GeV/nucleon has been successfully reproduced both by partonic models [4] and by hydrodynamics (i.e. soft physics dominated) [5]. This suggests that the complex interplay between hard and soft processes may not be unambiguously resolved by a single hadron spectrum at $p_T \leq 3$ -4 GeV/c.

In this paper we present a systematic study of low p_T hadron spectra, using all available data from $\sqrt{s} = 17$ -20

GeV/nucleon on π , K, p and Λ to constrain the magnitude of collective radial expansion. The expansion parameters so obtained are used to extrapolate the pion spectrum to high p_T for each collision system. The distribution is then compared to the π^0 data in order to determine at what p_T hydrodynamical expansion no longer accounts for the particle yield. As initial conditions and the surface profile of the expanding collision volume are not well constrained by the data, we choose the simplest profiles allowed by the data, and constrain the parameters using multiple hadronic species at transverse mass $m_T (= \sqrt{p_T^2 + m_0^2}) - m_0 \leq 1$ GeV. We include p+A collisions as well, because the secondary hadronic scattering and resulting transverse collective flow should be small.

Collective transverse flow boosts the velocity of the hadrons and introduces a hadron mass dependence to the slope of the m_T spectra. At CERN, NA44 [1] and NA49 [6] observed mass dependent inverse slopes (T_{eff}) of the m_T spectra, and interpreted the trend as common production from an expanding hadron gas. The data were fitted with a collective transverse flow velocity and temperature of the gas at freeze-out, when the hadrons cease to undergo strong interactions. The hadron spectra are not, however, completely exponential in m_T over a large range, and the deviation causes the extracted velocity and temperature parameters to depend upon the m_T fit range.

We select published effective temperatures that were determined over a common fit range for all the data, $(m_T - m_0) < 1.2$ GeV, by an exponential fit (Equation 1) to the measured m_T distributions for each hadron spectrum produced at midrapidity.

$$\frac{dN}{m_T dm_T} = A e^{-m_T/T_{eff}} \quad (1)$$

The experiments include NA44 [1, 7, 8, 9] and WA97 [10, 11] at the SPS; and, at the ISR, Alper et al. [12] and Guettler et al. [13]. For pions, the low- p_T region of $(m_T - m_0) < 0.3$ GeV, populated by decay of baryonic

resonances, was systematically excluded. The effective temperatures used are given in Table I with the references noted accordingly.

For all but p+p and p+Be, the π , K, p and Λ spectra were analyzed. The mass dependence of T_{eff} for positive particles is shown in Figure 1; it is clear that a linear increase with particle mass exists for heavy colliding systems.




FIG. 1: The effective temperatures are plotted against particle mass for positive particles produced in symmetric (A + A) collisions (left) and asymmetric (p + A and A + B) collisions (right) at CERN energies. Error bars are statistical only. In p+Pb (solid points), the effective temperature shows a slight dependence on particle mass.

The dependence of T_{eff} on both mass and the number of participating nucleons in the collision indicates radial expansion. The effective temperature includes the local temperature of a small piece of matter and its collective velocity (for motion in two dimensions, the classical $T = m_0 \langle v_x \rangle^2 / 2 + m_0 \langle v_y \rangle^2 / 2 = m_0 \langle v \rangle^2$, where $\langle v \rangle = \langle v_x \rangle = \langle v_y \rangle$ in circular coordinates in the plane transverse to the collision axis). Simple exponential fits treat each particle spectrum as a thermal source, so the collective expansion velocity cannot be extracted reliably from a single particle spectrum; however, by using the information from all the particles, the expansion velocity can be inferred. For example, NA44 [1] used

$$T_{eff} = T_{fo} + m_0 \langle \beta_t \rangle^2 \quad (2)$$

to separate the effective temperature into the two respective contributions (thermal and collective motion).

We fit equation 2 to the effective temperatures for particles produced in each collision system to determine the conditions at freeze-out. Equation 2 assumes that the flow velocity $\langle \beta_t \rangle$ is independent of particle mass m_0 (all particles collectively travel at the same radial velocity) [14]. It also assumes that the pions, kaons, and protons have the same freeze-out temperature T_{fo} .

Positive and negative hadrons yield equivalent freeze-out parameters. Omitting Λ and $\bar{\Lambda}$ from the fits with

Equation 2 does not change the resulting parameters. We use slopes from exponential fits to the measured m_T spectra, as these describe the data well over the limited m_T range $0.3 \leq m_T - m_0 < 1.2$ GeV. The simple relationship between T_{eff} , T_{fo} , and β_t was shown to agree well with hydrodynamical fits to the same data [1], justifying the use of this simple form.




FIG. 2: The freeze-out temperature T_{fo} (top) and mean radial expansion velocity $\langle \beta_t \rangle$ (bottom) plotted for each system in terms of system size AB at CERN energies. The open symbols are p + A systems which follow the trend of increasing collective flow with system size.

Figure 2 shows the dependence of the parameters on the size of the colliding system. The parameter T_{fo} remains approximately 140 MeV, regardless of system size; however, $\langle \beta_t \rangle$ increases, as would be expected from the larger pressure generated by copious particle production in the heavier colliding systems. Our values of T_{fo} and $\langle \beta_t \rangle$ are in reasonable agreement with other extractions of these parameters using two-particle correlations as well as single particle spectra. It should be noted that Figure 2 includes p+A results as well as p+p and A+A collisions (symmetric and asymmetric). The fact that p+Pb follows the trend of higher $\langle \beta_t \rangle$ for larger systems is remarkable, as significant collective flow in such collisions would not be expected. The effective temperatures most likely reflect the presence of initial state multiple parton scattering, often referred to as Cronin enhancement [15] already at relatively low hadron transverse momenta ($p_T \approx 1.3$ GeV/c for pions and 1.9 GeV/c for protons).

In heavy ion collisions, initial state multiple parton scattering should also be present [16]. At the relatively low $(m_T - m_0) < 1.2$ GeV fitted here, we observe a much larger mass dependence of the slopes in A+A than in

TABLE I: Inverse slope parameters (in MeV) of hadrons for p+p, p+nucleus, S+S, S+Pb, and Pb+Pb colliding systems at CERN energies. The errors are statistical and systematic, respectively.

Hadron	Pb+Pb	S+Pb	S+S	p+Pb	p+S	p+Be	p+p
π^+	$156 \pm 6 \pm 23^a$	$165 \pm 9 \pm 10^b$	$148 \pm 4 \pm 22^a$	$145 \pm 3 \pm 10^b$	$139 \pm 3 \pm 10^b$	$148 \pm 3 \pm 10^b$	$139 \pm 13 \pm 21^c$
K^+	$234 \pm 6 \pm 12^a$	$181 \pm 8 \pm 10^b$	$180 \pm 8 \pm 9^a$	$172 \pm 9 \pm 10^b$	$163 \pm 14 \pm 10^b$	$154 \pm 8 \pm 10^b$	$139 \pm 15 \pm 7^c$
p	$289 \pm 7 \pm 14^d$	$256 \pm 4 \pm 10^e$	$208 \pm 8 \pm 10^a$	$203 \pm 6 \pm 10^e$	$175 \pm 30 \pm 10^e$	$156 \pm 4 \pm 10^e$	$148 \pm 20 \pm 7^c$
Λ	$289 \pm 8 \pm 29^f$	—	—	$203 \pm 9 \pm 20^g$	—	—	—
$\bar{\Lambda}$	$287 \pm 13 \pm 29^f$	—	—	$180 \pm 15 \pm 18^g$	—	—	—

^aReference [1] (NA44 Collaboration).

^bReference [7] (NA44 Collaboration).

^cReference [12, 13] (ISR).

^dReference [9] (NA44 Collaboration).

^eReference [8] (NA44 Collaboration).

^fReference [10] (WA97 Collaboration).

^gReference [11] (WA97 Collaboration).

p+A collisions. In this soft momentum range, one would expect the boost from collective expansion to dominate over multiple scattering at the parton level. At higher m_T , the relative contributions should be reversed. This can be checked by attributing all the velocity boost to collective flow, extrapolating the spectrum to higher momentum under this assumption, and comparing to data at higher p_T or m_T .

In order to do this extrapolation, we assume that the soft physics processes in heavy ion collisions can be described by hydrodynamics. The parameters in Figure 2 are used to extrapolate to higher transverse momentum the spectrum of pions from a hydrodynamically expanding gas. We use a hydrodynamics parameterization from U. Heinz and coworkers, which assumes that all the particles decouple at a common freeze-out temperature T_{fo} , and that the source has an infinite length in the longitudinal direction (boost-invariance) [17].

The transverse velocity profile of the source is parameterized as a function of the radius $\xi = \frac{r}{R}$, where R is the maximum radius of the expanding source at freeze-out,

$$\beta_t(\xi) = \beta_s \xi^n. \quad (3)$$

β_s is the maximum surface velocity, and n (= 1 for linear) describes the profile. A flat particle density distribution $f(\xi) = \text{const}$ is also assumed. The geometrical average of the transverse velocity is therefore given by

$$\langle \beta_t \rangle = \frac{\int \beta_s \xi^n \xi d\xi}{\int \xi d\xi} = \frac{2}{2+n} \beta_s. \quad (4)$$

Different velocity profiles, such as n = 1/2 or 2 result in lower or higher surface velocities, respectively. If a different particle density distribution (for instance, a Gaussian) is used, then the average should be determined after weighting accordingly [18]. Using a linear velocity profile, the geometrical average flow velocity in the transverse plane is $\langle \beta_t \rangle = 2\beta_t/3$.

Each locally thermalized fluid element is given a transverse kick ρ that depends on the radial position

$$\rho = \tanh^{-1}(\beta_t(\xi)). \quad (5)$$

The transverse mass spectrum $\frac{dN}{m_T dm_T}$, of the particles is

$$A \int_0^1 m_t f(\xi) K_1\left(\frac{m_t \cosh(\rho)}{T_{fo}}\right) I_0\left(\frac{p_t \sinh(\rho)}{T_{fo}}\right) \xi d\xi \quad (6)$$

which we integrate numerically.

We use Equation 6 with the freeze-out temperature T_{fo} and the maximum surface velocity β_t from Figure 2 to extrapolate the particle spectra to high p_T . The normalization A is allowed to vary, as we compare only the shape of the extrapolated distribution to data. Figure 3 shows the resulting extrapolated spectra compared to the neutral pion spectra in Pb+Pb, S+Pb, S+S [19, 20] and charged pions in p+W collisions [15]. (The equivalent p+Pb pion data in this momentum range was not available.) The dashed lines are error bands, corresponding to uncertainties in T_{fo} and β_t from statistical and systematic errors in the effective temperatures for each collision system.

For heavy ion collisions, the measured π^0 p_T spectra are consistent with hydrodynamic distributions, though the highest p_T point tends toward the edge of the upper error band. The moderate p_T values measured to date in heavy ion collisions contain a mix of hard and soft processes. To cleanly observe hard processes in heavy ion collisions, measurements at $p_T \geq 4$ GeV/c will be very valuable.

Pions produced in p+W collisions clearly deviate from the hydrodynamic parameterization at $p_T > 2$ GeV/c, underscoring the differences in the physics between p+A and A+A collisions. The long tail in the p+W spectrum presumably reflects initial state parton multiple scattering and the onset of hard processes, which are masked by the contribution of boosted soft particles in heavy ion




FIG. 3: The hydrodynamically extrapolated pion spectrum compared to data to the highest available p_T for multiple collision systems at CERN. The dashed lines indicate error bands for each boosted spectrum. The hydrodynamic distribution clearly deviates from the p+W data at $p_T > 2$ GeV/c, while no such discrepancy is observed in the heavier colliding nuclei, illustrating the physics differences between p+A and A+A.

collisions. By extrapolation, the p+A data may be used to estimate the contribution of multiple scattering of partons to the observed broadening of the hadron spectra in A+A collisions.

Stimulating discussions with Vladislav Pantuev, Xin-Nian Wang, Nu Xu, Axel Drees, R. Soltz, and S.C. Johnson are gratefully acknowledged. This work was supported by the U.S. Department of Energy under grant No. DE-FG02-96ER40988.

[1] I.G. Bearden et al. (NA44 Collaboration), Phys. Rev. Lett. **78**, 2080 (1997).
[2] H. Appelshauser et al (NA49 Collaboration), Eur. Phys. J. **C2**, 661 (1998).
[3] X.N. Wang, Phys. Rept. **280**, 287 (1997).
[4] X.N. Wang, Phys. Rev. Lett. **81**, 2655 (1998).
[5] M.M. Aggarwal et al (WA98 Collaboration), Phys. Rev. Lett. **83**, 926 (1999).
[6] H. van Hecke, H. Sorge and N. Xu, Phys. Rev. Lett. **81**,

5764 (1998).
[7] H. Boggild et al. (NA44 Collaboration), Phys. Rev. C **59**, 328 (1999).
[8] I.G. Bearden et al. (NA44 Collaboration), Phys. Rev. C **57**, 837 (1998).
[9] I. Bearden et al., Phys. Lett. B **388**, 431 (1996).
[10] E. Andersen et al., Journ. Phys. G **25**, 171 (1999).
[11] WA97 Collaboration, Phys. Lett. B **433**, 209 (1998).
[12] B. Alper et al., Nucl. Phys. B **100**, 237 (1975).
[13] K. Guettler et al., Nucl. Phys. B **116**, 77 (1976).
[14] T. Csorgo and B. Lorstad, Phys. Rev. C, **54**, 1390, (1996).
[15] J.W. Cronin et al., Phys. Rev. D. **11**, 3105 (1975).
[16] E. Wang and X.-N. Wang, Phys. Rev. C **64**, 034901 (2001).
[17] E. Schnedermann, J. Sollfrank and U. Heinz, Phys. Rev. C **48**, 2462 (1993).
[18] S. Esumi, S. Chapman, H. van Hecke and N. Xu, Phys. Rev. C **55**, R2163 (1997).
[19] M.M. Aggarwal et al (WA98 Collaboration), Phys. Rev. Lett. **81**, 4097 (1998).
[20] R. Albrecht et al (WA80 Collaboration), Eur. Phys. J. **C5**, 255 (1998).